PCI

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵: C12N 15/00, 15/87, 15/23 A01K 67/027, C12N 5/10

A1 (11) International Publication Number:

WO 93/24626

(43) International Publication Date:

9 December 1993 (09.12.93)

(21) International Application Number:

PCT/US92/04426

(22) International Filing Date:

28 May 1992 (28.05.92)

(71) Applicant (for all designated States except US): SCIENTIF-IC DIMENSIONS USA, INC. [US/US]; 185 Alewife Brook Parkway, Cambridge, MA 02138 (US).

(72) Inventors; and

7

(75) Inventors, and
(75) Inventors/Applicants (for US only): LARIONOV, Oleg
[RU/RU]; DOBROVOLSKY, Vasily N. [RU/RU];
Shemyakin Institute of Bioorganic Chemistry, Russian
Academy of Sciences, ul. Miklukho-Maklaya, 16/10,
GSP Moscow V-437, 117871 (RU).


(74) Agents: GREASON, Edward, W. et al.; Kenyon & Kenyon, One Broadway, New York, NY 10004 (US).

(81) Designated States: AT, AU, BB, BG, BR, CA, CH, CS, DE, DK, ES, FI, GB, HU, JP, KP, KR, LK, LU, MG, MN, MW, NL, NO, PL, RO, RU, SD, SE, UA, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LU, MC, NL, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, SN, TD, TG).

Published

With international search report.

(54) Title: TRANSGENIC ANIMAL PRODUCTION WITH BIOLISTICALLY TRANSFORMED SPERMATOZOA



(57) Abstract

A method of producing transgenic animals which comprises binding DNA which expresses a desired trait to a particle, inserting the carrier particle with bound DNA into spermatozoa, and fertilizing a sufficiently related egg. Also disclosed is a method for reversibly fixing spermatozoa on a slightly dried agarose gel so that they can be bombarded by DNA-laden particles from a gene gun.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
ΑU	Australia	GA	Gabon	MW	Malawi
8B	Barhados	GB	United Kingdom	NL	Netherlands
BE	Betgium	GN	Guinea	NO "	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BC	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	1E	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic	SD	Sudan
CC	Congo		of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SK	Slovak Republic
CI	Côte d'Ivoire	ΚZ	Kazakhstan	SN	Senegal
CM	Cameroon	L.)	Liechtenstein	S U	Soviet Union
CS	Czechoslovakia	LK	Sri Lanka	TD	Chad
CZ	Czech Republic	I.U	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	UA	Ukraine
DK	Denmark	MC	Madagascar	US	United States of America
ES	Spain	Ml.	Mali	VN	Viet Nam
FI	Finland	MN	Mongolia		

TRANSGENIC ANIMAL PRODUCTION WITH BIOLISTICALLY TRANSFORMED SPERMATOZOA

FIELD OF THE INVENTION

The present invention relates to methods of creating transgenic animals which contain cloned genes taken up into spermatozoa such that the spermatozoa function as vectors when fertilizing an egg of a sufficiently related species. The spermatozoa take up the cloned DNA, for example, as a result of being bombarded by DNA-laden particles from a gene gun while fixed to an agarose gel. The cloned sequence is transmittable to offspring.

BACKGROUND OF THE INVENTION

Introducing foreign DNA into somatic and germinal cells of animals is well established [Palmiter & Brinsten, Ann. Rev. Genet. 20, 465-499 (1986)]. The most widely used method is the direct microinjection of foreign DNA into pronuclei of fertilized eggs [Gordon et al., Pros. Natl. Acad. Sci. USA 77, 7380-84 (1980)]. However, receiving zygotes at the developmental (two pronuclei) stage suitable for microinjection is problematic. Introducing foreign DNA into mammalian genome may also be done using transfected embryo-derived stem (ES) cells, [Lovell-Badge et al., Cold Spring Harb. Symp. Quant. Biol. 50, 707-11 (1985)], although obtaining live-stock ES cell lines is extremely difficult. The insertion of genes into the germ line via retroviral vectors has also been improved [van der Putten, Pros. Natl. Acad. Sci. USA 82, 6148-52 (1985)]. Nonetheless, because of the structural characteristics of retroviruses, using them to construct vectors presents grave difficulties.

Only recently has cloned DNA been introduced directly into spermatozoa [Lavitrano et al., Cell <u>57</u>, 717-723 (1989)]. Indeed, mature

2

mouse sperm cells incubated in an isotonic buffer with cloned DNA will capture DNA molecules. Thus, sperm cells may be used as vectors for introducing foreign DNA into eggs at fertilization. Mouse eggs can be fertilized in vitro by sperm incubated in the cloned DNA solution then 5 transferred, at the two-cell stage, to the oviducts of foster mothers. Approximately 30% of 250 screened mice revealed clear sequence homology to the cloned DNA sequence. The key to this method is the capture of foreign DNA by spermatozoa through exposure in an isotonic solution.

Foreign genes have also been introduced into spermatozoa using liposomes, [Hofer, Europ. Biotech. Newsletter 47, 4704 (1988)]. The 10 spermatozoa loaded by the liposome method have been used to transform egg or somatic cells under intra- or extracorporal conditions to produce hepatitis surface antigen and a rat immunoglobulin. [German Pat. No. WO 87/05325].

15

Transgenic animals also have been produced by introducing foreign DNA into some tissues of adult animals or fetuses by means of: a) a retrovirus vector carrying the DNA of interest; b) CaPO₄ mixed with the DNA of interest, or c) electroporation, [U.K. Patent No. 2,223,755]. Unfortunately, use of these treatments results in animals which do not transmit foreign DNA to their progeny. Finally, a transgenic non-human 20 mammal has been produced where germ cells and somatic cells contain a recombinant activated oncogene sequence introduced into said mammal at an embryonic stage, [U.S.P. 4,736,866 (The "Harvard Transgenic Mouse")].

"Gene guns" have been manufactured to facilitate introduction of cloned DNA into various types of cells by bombarding them. One typical example of a gene gun, Dupont's PDS-2000, utilizes microparticles (usually made of tungsten or gold) laden with the cloned DNA to pierce a cell membrane and introduce the DNA into the cell. Specifically, Dupont's PDS-2000 has been used on muscle cells, human cultured cells, fibroblasts, chinese 30 hamster ovaries and lymphocytes to cause the up-take of cloned DNA. However, the gene gun requires its target cells to be exposed on a support

3

surface to prevent particle breaking in the medium and for efficient DNA placement. The initial conditions for using Dupont's PDS-2000 vary, but Dupont suggests the following conditions to optimize bombardment: cells should be maintained at 15 inches of Hg, for as brief a period as possible (1-3 min.) optimizing vacuum pressure up to 27 inches of Hg in three such integrals. This increases the speed of the microcarriers. If cells become damaged or cell survival decreases, vacuum levels should be reduced. In addition, as the distance between the accelerator and the stopping plate increases, the velocity of the microcarrier decreases, and dispersion increases. Thus, different ranges of distance should be tried for comparison. The microcarrier must be placed optimally in the acceleration tube so the velocity is optimized. Microcarriers should be one-tenth the diameter of the cell. M5

(average is 0.35 μm) and M10 (average is 0.73 μm) tungsten and the one μm gold particles have all been used successfully on animal cells in tissue culture
 [DuPont Biolistic™ PDS-1000 Instruction Manual]. The successful bombardment of live sperm has not been yet reported.

DESCRIPTION OF THE FIGURES

Figure 1 is a schematic diagram of a recombinant vector used to transfer sperm in a preferred embodiment of the present invention.

Figure 2 summarizes the various general steps of certain preferred embodiments of the invention.

Figure 3 is a sketch of the vector construction used on the BLG gene for human gamma interferon expressing in mammary gland of transgenic animals.

25

SUMMARY OF THE INVENTION

The purpose of this invention is to create a universal, simple, efficient and inexpensive method for production of transgenic live stock transmitting foreign genes to their progeny. This method will obviate the difficulties

4

inherent with recovering early embryos and treating donor and acceptor animals.

In accordance with the present invention, a method of producing a transgenic animal is disclosed where DNA which expresses a desired trait is bound to a carrier particle, the carrier particle is inserted into spermatozoa, and the spermatozoa is used to fertilize an unfertilized egg from a species sufficiently related to the one from which said spermatozoa was collected. Species from which spermatozoa and an unfertilized egg are collected are "sufficiently related" if fertilization of the egg by the sperm can result in a viable embryo. In certain preferred embodiments, a gene gun can be used to insert a carrier particle bound with DNA into spermatozoa. Any method of inserting the DNA-laden carrier particle can be used, although bombardment with a gene gun is preferred.

The carrier particles to which the cloned DNA is bound are tungsten or gold. Any other material may be used to bind the cloned DNA if it will sufficiently survive the chosen insertion method and properly present the DNA for integration into the spermatozoan genome.

Fertilization of the unfertilized egg by the spermatozoa containing the inserted DNA can be by *in vitro* fertilization, by artificial insemination (for example, in the oviduct), or by any other means for uniting genetic material from gametes.

A method is also disclosed for reversibly fixing sperm so that they may be successfully bombarded by DNA-laden particles. Sperm recovered from epididymis is dissolved at room temperature in a small volume of medium, then dropped onto the surface of slightly dried 1% agarose gel. After liquid evaporation, the spermatozoa become fixed on the gel. The bombardment process can occur for three to five minutes, after which a medium is added to the gel so the spermatozoa can regain motility.

DETAILED DESCRIPTION OF THE INVENTION

This invention, in certain preferred embodiments, consists of the following steps (see fig. 2): 1) fixing spermatozoa onto agarose gel, 2) bombarding fixed spermatozoa (head dimensions are approximately $2x5\mu m$) 5 by DNA-coated particles using a gene gun (gold or tungsten particles with diameter of $0.2\text{-}0.5\mu m$); 3) harvesting bombarded spermatozoa and using centrifugation in a Percoll density gradient to isolate spermatozoa containing DNA-coated particles; and 4) using isolated spermatozoa for artificial insemination. After fertilizing an egg with spermatozoa carrying DNA-coated particles, a male pronucleus will form, followed by dissociation of the foreign DNA from the particle surface and integration of it into the genome. The particle will be torn away during embryo cleavage.

reaction (PCR)] of the 3'-end of the beta-lactoglobulin (BLG) gene

comprising the last intron and poly(A) site and inserting of that fragment into a Pvu II site (fig. 1). More specifically, the 11.3kb DNA fragment comprising the 4.9kb transcription unit, the 4.8kb 5' flanking sequence and the 1.6kb 3' flanking sequence of the BLG gene was cloned (using standard methods of genetic engineering [Davis et al., Basic methods in Molecular Biology.

Elsivier, N.Y. (1986)] into plasmid pBRUC318 (fig. 3.A) from a sheep genomic library (in phage lambda-EMBL3) using as probes two partially overlapping synthetic oligodeoxyribonucleotides Seq. Id. No:1 and Seq. Id. No:2 complementary to the coding sequence of the BLG gene first exon [Ali S., Clark, A.J., J. Mol. Biol. 199, 415-426 (1988)]. The PvuII site resides within the first exon (in untranslated region). Human gamma-interferon

A vector was constructed by amplifying [using polymerase chain

A vector was created for human gamma-interferon (g-IFN) expression in the mammary gland. A fragment of the 3' end of the BLG gene, comprising the last intron and poly(A) site, was amplified using polymerase chain reaction (PCR) (fig.3.B.). For this purpose two synthetic oligonucleotides Seq. Id. No:3 and Seq. Id. No:4 were used as primers. They

cDNA was inserted into the SnaBI site.

6

contained recognition sequences for restriction endonucleases SnaBI and Hind III at their 5' ends and were complementary to flanks of this fragment. PCR conditions were: 100pmol of each primer, 0.1ng of target DNA (BLG gene), 67mM TrisHCl pH8.8, 16.6mM (NH₄)₂SO₄, 6.7mM MgCl₂, 10mM DTT, 0.2mM of each dNTP, 100µg/ml BSA, Taq I DNA polymerase 2un, H₂O to final volume of 100µl. A total of 30 cycles were performed, each cycle being 30 sec. at 95°C, 30 sec. at 65°C, 2 min. at 72°C. An amplified fragment (616bp) bearing newly created SnaBI (at its 5' end) and Hind III (at its 3' end) sites was ligated into the Pvu II site that resides within the untranslated region of the BLG gene first exon using standard methods of molecular cloning (fig.3.C). g-IFN cDNA [received from Prof. E.D. Sverdlov (Institute of Bioorganic Chemistry, USSR Academy of Sciences)] that contained initiating and terminating codons was inserted into the SnaBI site (fig.3.D). This DNA construct is used to load carrier particles.

DNA is bound to tungsten or gold particles using the following procedure: 1.25 mg of tungsten is heated in 95% EtOH at 65°C for four hours. 25 μ g of DNA, 2.5M CaCl₂ and 0.1M spermidine are combined to a final volume of 575 μ l (final concentration of 1.1M CaCl₂ and 8.7mM spermidine). The mixture is vortexed at 4°C for 10 minutes, then centrifuged at a low speed (500 rpm) for 5 minutes. Approximately 550 μ l of supernatant is then removed and the remaining 25 μ l is separated into 1 μ l aliquots for loading onto a macrocarrier. It is better to use gold particles as tungsten may be toxic. DNA is bound to gold particles according to the method described by Du Pont. [Newsletter, A Publication of the Du Pont Company, 1, 1-4 (1990)].

15

20

25

To create transgenic animals, spermatozoa are bombarded by accelerated DNA-coated microparticles from a gene gun, such as the PDS-2000. For the purpose of bombardment, spermatozoa are reversibly fixated on a low melting point agarose gel. To accomplish the reversible fixation, sperm is collected from the epididymis, and dissolved at room temperature in a small volume (approx. 50-150µl) of M2 medium [Quinn, et al., J. Reprod.

7

Fertil. 66, 161-168 (1982). The sperm solution is then dropped onto the surface of a slightly dried 1% agarose gel (Sea Plaque agarose, FMC, USA). After liquid evaporation, the spermatozoa are fixed on the gel. Three to five minutes later, during which time bombardment with foreign DNA can occur, M2 medium $(50-250\mu l)$ is added to the gel to reverse fixation. About 60-80% of fixed spermatozoa regain motility in the liquid phase.

Spermatozoa containing DNA-coated particles are separated from other spermatozoa by density gradient centrifugation. The density gradient is formed in 10-15ml centrifuge tubes during high-speed centrifugation (15 min. 29000g) of Duldecco's PBS containing 60-90% of Percoll. Bombarded spermatozoa washed over the surface of the gel are layered onto preformed density gradient tubes and after low-speed centrifugation (20 min. 400g) at room temperature, they are separated on fractions depending on their floating density. Spermatozoa which contain particles are higher density so they concentrate closer to the bottom of the tube. The desired spermatozoa are then collected by puncturing of the tube bottom.

The DNA-laden spermatozoa are then used to artificially inseminate a female specimen or to fertilize eggs by other known means. For example, with mice the preferred method of fertilization is to introduce DNA-laden spermatozoa directly into the oviduct of pseudopregnant females (females mated by sterile males) or transfer the selected spermatozoa (from density gradient tube) to the the oviduct or the uterus using fine glass pipet (outer diameter 100-120µm) This procedure is analogous to the procedure of embryo transfer into the uterus [Hogan et al., Manipulating the Mouse Embryo. (Cold Spring Harbor Laboratory, N.Y., 142-145 (1986)]). The normal artificial insemination of a mouse female is difficult due to particularities of organization of mouse female reproductive organs. In vitro fertilization is also possible.

For livestock routine artificial insemination is preferred, using the same gene gun technology applied with mouse spermatozoa. In vitro fertilization is also possible.

8

Integration of foreign DNA into the murine (or other) genome is analyzed using PCR. Two oligonucleotides (e.g., Seq. Id. No:5 and Seq. Id. No:6) are used as primers for checking the integration of the g-IFN cDNA. They are complementary to g-IFN cDNA and form 252bp DNA fragments during PCR. PCR conditions: 20pmol of each primer, 0.1-0.5µg of mouse genomic DNA, 0.2mM of each primer, 50mM KCl, 10mM TrisHCl pH8.14, 1.5mM MgCl₂, 100µg/ml gelatin, Taq I DNA polymerase 1-1.5 units, and H₂0 to final volume of 20µl. The mixture runs for 30 cycles, each being: 30sec 94°C, 30sec 55°C, 30sec 72°C. 10-15µl aliquots of reaction mix is electrophoresed on 1.5% agarose gel in the presence of ethidium bromide at 100V for 30min and viewed under UV light.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: Larionov, Oleg Dobrovolsky, Vasily N.

(ii) TITLE OF INVENTION: A NEW METHOD FOR TRANSGENIC ANIMAL PRODUCTION

(111) NUMBER OF SEQUENCES: 6

CORRESPONDENCE ADDRESS: (iv)

(A) ADDRESSEE: Paul Lempel, Esq. (B)

STREET: One Broadway

CITY: New York

STATE: New York COUNTRY: USA

ZIP: 10004 0000 3

COMPUTER READABLE FORM:
(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS

SOFTWARE: Patentin Release #1.0, Version #1.25

00

(A) APPLICATION NUMBER: US N/A CURRENT APPLICATION DATA: (vi)

FILING DATE: 16-SEP-1991 CLASSIFICATION:

E 0

ATTORNEY/AGENT INFORMATION: (viii)

(A) NAME: Lempel Esq., Paul(B) REGISTRATION NUMBER: 21,198(C) REFERENCE/DOCKET NUMBER: 1036/61

(A) TELEPHONE: (212) 425-7200 TELECOMMUNICATION INFORMATION: (ix)

10

23

56

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

(2) INFORMATION FOR SEQ ID NO:2:

(i)

GIGGCGICCA GGCCATCAIC GIC

SEQUENCE CHARACTERISTICS:
(A) LENGTH: 26 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

TTTCATGGTC TGGGTGACGA TGATGG

(B) TELEFAX: (212) 425-5288

(2) INFORMATION FOR SEQ ID NO:1:

SEQUENCE CHARACTERISTICS:
(A) LENGTH: 23 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear (ï)

LENGTH: 23 base pairs

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

11

25

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

(A) LENGTH: 25 base pairs (B) TYPE: nucleic acid (C) STRANDEDNESS: single (D) TOPOLOGY: linear

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

SEQUENCE CHARACTERISTICS:

(i)

(2) INFORMATION FOR SEQ ID NO:3:

TACGTAGGTG AGCCCCTGCC GGTGC

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25 base pairs (B) TYPE: nucleic acid (C) STRANDEDNESS: single (D) TOPOLOGY: linear

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

AAGCTTCCAG CAAAGACTCA GAAGG

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

LENGTH: 25 base pairs TYPE: nucleic acid STRANDEDNESS: single TOPOLOGY: linear

€£0£

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

12

25

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

AATGCAGGTC ATTCAGATGT AGCGG

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 25 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

CGAATAATTA GTCAGCTTTT CGAAG

13

CLAIMS

We claim: 1 1. A method of producing a transgenic animal said method 2 comprising: 3 A. binding DNA which expresses a desired trait to a carrier 4 particle; 5 В. inserting said carrier particle with said DNA into 6 spermatozoa of a selected species; and 7 fertilizing an unfertilized egg from a species sufficiently 8 related to said selected species. 1 2. The method of claim 1 wherein a gene gun is used to insert 2 said carrier particle bound with DNA into said spermatozoa. 1 3. The method of claim 2 wherein said carrier particle is made of 2 a material selected from the group consisting of tungsten and gold. 1 4. The method of claim 1 wherein said unfertilized egg is fertilized 2 in vitro. 1 5. The method of claim 1 wherein said unfertilized egg is fertilized 2 by artificial insemination. 6. The method of claim 1 further comprising separating 1 2 spermatozoa containing said carrier particle from spermatozoa not carrying 3 said carrier particle. . 1 7. The method of claim 6 wherein said spermatozoa are separated 2 by centrifugation.

WO 93/24626

1	8.	The method of claim 2 further comprising, prior to inserting
2	said carrier p	particle, reversibly fixing said spermatozoa to a support.
	_	
1	9.	The method of claim 8 wherein said spermatozoa are fixed by a
2	process acco	rding to claim 10.
1	10.	A method for reversibly fixing spermatozoa said method
2	comprising:	
3		(a) diluting a sample of spermatozoa in a first medium;
4		(b) contacting said sample with an agarose gel; and
5		(c) evaporating the liquid portion of said first medium such
6	that said spe	rmatozoa become fixed to said gel.
1	11.	The method of claim 10 further comprising reversing the
2	fixation of sa	aid sperm by adding a second liquid medium to said spermatozoa.
1	12.	The method of claim 10 which the first medium is a M2
2	medium.	
1	13.	The method of claim 10 wherein the second medium is an M2
2	medium.	The method of claim to wherein the second medium is an M2
4	medium.	
1	14.	The method of claim 10 wherein said gel is 1% agarose gel.
1	15.	A method for reversibly fixing spermatozoa comprising:
2	collecting sp	erm from the epididymis and dissolving it at room temperature in
3	a small volu	me (approx. 50-150µl) of M2 medium; dropping the sperm
4	solution ont	o the surface of a slightly dried 1% agarose gel; allowing time for
5	liquid evapo	oration so that the spermatozoa become fixed on the gel;
6	hombarding	the fixed spermatozoa for three to five minutes; and adding M2

7 medium (50-250 μ l) to the gel to reverse fixation.

6

method of claim 11.

15

16. 1 The method of claim 15 wherein reversibly fixed spermatozoa are fixed according to the method of claim 10. 1 17. A DNA-laden spermatozoon made according to the method of claim 18. 2 1 18. A method of production of DNA-laden spermatozoa 2 comprising: the method for reversibly fixing spermatozoa according 3 (a) to the method of claim 10; and 4 5 the method for reversing the fixation according to the (b)

1/3




FIG. 1

2/3




FIG. 2

SUBSTITUTE SHEET




FIG. 3 SUBSTITUTE SHEET

International Application No

		ECT MATTER (if several classification :				
	to International Patent 5 C12N15/00 C12N5/10			A01K67/027		
II. FIELDS	SEARCHED					
		Minimum Docum	entation Searched ⁷			
Classificati	Classification System Classification Symbols					
Int.Cl.	. 5	C12N ; C07K				
			than Minimum Documentation are Included in the Fields Searched ⁸			
		ED TO BE RELEVANT ⁹				
Category °	Citation of Do	ocument, 11 with indication, where appropr	iate, of the relevant passages 12	Relevant to Claim No. ¹³		
x	26 Decen	119 781 (AGRACETUS) mber 1991 e 6, line 25 - page 12,	. line 30	1-3		
x	WO,A,8 7 11 Septe	705 325 (TRANSGENE) ember 1987 whole document		1		
Y	12 June	431 839 (SCOPUS-GENETION 1991 whole document	CS)	1-5		
Y	RICERCHE 26 July	008 192 (CONSIGLIO NAZI E) 1990 whole document	ONALE DELLE	1-5		
Y		 118 991 (E. I. DUPONT [mber 1991	DE NEMOURS)	1-3		
			-/			
"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "C" document published prior to the international filing date but later than the priority date claimed						
IV. CERTIF						
Date of the	Actual Completion of t	the International Search ARY 1993	Date of Mailing of this International S	earch Report		
International	Searching Authority	AN PATENT OFFICE	Signature of Authorized Officer CHAMBONNET F.J.			

III. DOCUM	ENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)	
Category °	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
Υ	WO,A,9 107 487 (DUKE UNIVERSITY) 30 May 1991 see the whole document	1-5
х	WO,A,9 100 359 (AGRACETUS) 10 January 1991 see page 5, line 10	1-3
Y	see the whole document	1-5
х	WO,A,9 003 439 (AMGEN) 5 April 1990 see the whole document	1
X	WO,A,9 008 832 (NATIONAL RESEARCH DEVLOPMENT CORPORATION) 9 August 1990 see the whole document	1
		·
!		
	·	

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.

US 9204426 SA 60799

This armex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on

The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

20/01/93

Patent document cited in search report	Publication date	Patent family member(s)		Publication date	
WO-A-9119781	26-12-91	US-A- AU-A- CA-A- EP-A-	5149655 8210891 2065361 0487711	22-09-92 07-01-92 22-12-91 03-06-92	
WO-A-8705325	11-09-87	DE-A- AU-B- AU-A- EP-A- JP-T- ZA-A-	3636991 597965 7167987 0258427 63502638 8701468	24-09-87 14-06-90 28-09-87 09-03-88 06-10-88 17-08-87	
EP-A-0431839	12-06-91	AU-A- CA-A- JP-A-	6767290 2031344 3266983	04-07-91 04-06-91 27-11-91	
WO-A-9008192	26-07-90	AU-A- EP-A- JP-T-	4810790 0453458 4504352	13-08-90 30-10-91 06-08-92	
WO-A-9118991	12-12-91	AU-A-	7878891	31-12-91	
WO-A-9107487	30-05-91	AU-A- EP-A-	6964191 0500799	13-06-91 02-09-92	
WO-A-9100359	10-01-91	AU-A- CA-A- EP-A- JP-T-	5856790 2019676 0431135 4500314	17-01-91 26-12-90 12-06-91 23-01-92	
WO-A-9003439	05-04-90	US-A- AU-A- EP-A- JP-T-	5162215 4312389 0393176 3503485	10-11-92 18-04-90 24-10-90 08-08-91	
WO-A-9008832	09-08-90	AU-A- EP-A- GB-A,B JP-T-	5027690 0455695 2228487 4503004	24-08-90 13-11-91 29-08-90 04-06-92	

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.